



UDC 511.3

An Estimate of a Certain Summatory Functions Class

V. A. Matveev

Saratov State University, Russia, 410012, Saratov, Astrakhanskaya st., 83, vladimir.matveev@gmail.com

In this paper summatory functions of form $\sum_{n \leq x} h(n)n^{it}$, $1 \leq |t| \leq T$ for finite-valued functions $h(n)$ of natural argument with bounded sum function are estimated.

Key words: dirichlet character, summatory functions, power series.

References

1. Chudakov N. G., Bredikhin B. M. Application of Parseval's identity in estimations of summatory functions of Dirichlet characters on numerical semigroups. *Uspekhi matematicheskikh nauk*, 1956, vol. 9, pp. 347–360 (in Russian).
2. Dem'ianov V. F., Malozemov V. N. *Vvedenie v minimaks* [Introduction to minimax]. Moscow, Nauka, 1982 (in Russian).
3. Gel'fond A. O. Ob arifmeticheskom ekvivalente analitichnosti L -riada Dirikhle na priamoi [On certain arithmetical equivalent of analytic property of Dirichlet L -series on $\text{Re } s = 1$ line]. *Izbrannye trudy* [Selectas], Moscow, Nauka, 1973. pp. 310–328 (in Russian).
4. Matveev V. A., Matveeva O. A. On behavior in critical strip of Dirichlet series with finite-valued coefficients and bounded summatory function. *Chebyshevskii sbornik* [Chebyshev collection], 2012, vol. 13, iss. 2, pp. 106–116 (in Russian).